

STANDARD OPERATING PROCEDURES

DIVISION OF COMPARATIVE MEDICINE
UNIVERSITY OF SOUTH FLORIDA

SOP#: 427.2

Date Issued: 10/18

Date Revised: 12/23

Page 1 of 3

TITLE:	Food and Fluid Regulation in Mice and Rats
SCOPE:	All Program Personnel
RESPONSIBILITY:	Principal Investigators
PURPOSE:	To Outline the Proper Procedures to be Followed for Food and Fluid Regulation in Mice and Rats.

I. PURPOSE

1. This SOP outlines the proper procedures to be followed for food and fluid regulation in mice and rats.
2. Food and/or fluid regulation may be used in studies (a) to motivate animals to perform novel or learned tasks, (b) of the behaviors and mediators of hunger and thirst, and (c) of the homeostatic regulation of energy metabolism and its influence on disease pathogenesis.
3. This SOP will outline important issues to be considered and addressed in IACUC protocols that require food or fluid regulation.

II. RESPONSIBILITY

1. It is the responsibility of the Principal Investigator to ensure the research staff are properly trained, and follow both the IACUC protocol and this SOP.

III. BACKGROUND & SCOPE

1. The objective when food and fluid regulation studies are planned and executed should be to **use the least regulation necessary to achieve the scientific objectives while maintaining animal well-being**. In order to accomplish this objective, the following must be considered and described in the IACUC protocol:
 - a. The level of food and/or fluid regulation
 - b. Potential adverse consequences of the regulation
 - c. Methods to assess the health and well-being of animals under regulation
2. Definitions
 - a. **Ad libitum**: Rodents are offered access to a continuous supply of food and water.
 - b. **Deprivation**: Complete withholding of food or water.
 - c. **Fasting**: Acute (<12hr) withholding of food or water prior to imaging, behavior testing, etc.
 - d. **Regulation/Scheduling**: Controlling the **amount of, and when** water or food is offered to an animal; an amount less than the average daily *ad libitum* consumption.
3. A scientific **justification** must be provided in the IACUC protocol when proposing food and/or fluid regulation. This justification should provide a **rationale for the level and length of time** the regulation is proposed.


IV. PROCEDURES

1. Protocols requiring chronic (i.e., >24 hrs) regulation should indicate that either a strain-specific-published or in-house-determined ***ad libitum* intake normal range** will be used.
2. Protocols should provide the **formula used for the proposed regulation** (e.g. 70% of *ad libitum* intake food regulation to achieve 80% body weight of controls). Typically, food regulation should not be lower than 30% of *ad libitum* intake.
3. Note that acute water deprivation of up to 24 hours will result in clinical dehydration (e.g. listless, inactive, “skin tenting”, sunken eyes). Chronic **water regulation** of up to 50% of *ad libitum* offering may be imposed gradually over 7 days.
4. Consideration should be made to allow **food and water** to be available **concurrently**, as rodents typically do not eat without available water. During regulation, access to food and/or water must be for 15 minutes at a minimum.
5. Records must be kept of rodents on regulation, including the **baseline body weight**, records of **daily monitoring** ensuring the availability of food and water, at least **twice weekly body weights**, a **weekly body condition score** (BCS, see Appendix A), and other observations (e.g., hydration status).
6. Animal assessments must be made on **CMDC #245** entitled ***Food and Fluid Regulation Record***. Records must be kept in the housing room book while animals are present, and be made available for review.
7. If animals are undergoing Fasting, then documentation (CMDC #245) is not necessary at that time.
8. Specific **clinical endpoints** that require intervention must be clearly stated in the protocol (e.g., **food regulated rodent may not lose >20% of baseline body weight** without adjusting the daily food allowance) (e.g., **fluid regulated rodent may not lose >10% of baseline body weight** without adjusting the availability of fluid).
9. Rodents that appear dehydrated must immediately be provided with a measured volume of drinkable fluid *ad libitum*, and alternative fluid sources (e.g. hydrogel, sterile subcutaneous fluids).

Approved:

Date:

Body Condition Scoring (BCS) Charts for Mice and Rats



- BC 1**
Rat is emaciated
 - Segmentation of vertebral column prominent if not visible.
 - Little or no flesh cover over dorsal pelvis. Pins prominent if not visible.
 - Segmentation of caudal vertebrae prominent.
- BC 2**
Rat is under conditioned
 - Segmentation of vertebral column prominent.
 - Thin flesh cover over dorsal pelvis, little subcutaneous fat. Pins easily palpable.
 - Thin flesh cover over caudal vertebrae, segmentation palpable with slight pressure.
- BC 3**
Rat is well-conditioned
 - Segmentation of vertebral column easily palpable.
 - Moderate subcutaneous fat store over pelvis. Pins easily palpable with slight pressure.
 - Moderate fat store around tail base, caudal vertebrae may be palpable but not segmented.
- BC 4**
Rat is overconditioned
 - Segmentation of vertebral column palpable with slight pressure.
 - Thick subcutaneous fat store over dorsal pelvis. Pins of pelvis palpable with firm pressure.
 - Thick fat store over tail base, caudal vertebrae not palpable.
- BC 5**
Rat is obese
 - Segmentation of vertebral column palpable with firm pressure; may be a continuous column.
 - Thick subcutaneous fat store over dorsal pelvis. Pins of pelvis not palpable with firm pressure.
 - Thick fat store over tail base, caudal vertebrae not palpable.